arXiv:astro-ph/0512641v2 20 Jan 2006

Analytic Results for a Flat Universe Dominated by Dust and Dark Energy

A. Gruppuso, F. Finelli*
INAF/IASF-BO, Istituto di Astrofisica Spaziale e Fisica Cosmica, Sezione di Bologna
Istituto Nazionale di Astrofisica, via Gobetti 101, I-40129 Bologna - Italy[†]

(Dated: February 5, 2008)

We find the solution for the scale factor in a flat Universe driven by dust plus a component characterized by a constant parameter of state which dominates in the asymptotic future. We also present an analytic formula (in terms of hypergeometric functions) for the past light cone in such a universe. As applications of this result, we give analytic expressions for the Luminosity Distance and for the Acoustic Scale, where the latter determines the peaks positions in the pattern of CMB anisotropies.

PACS numbers: 98.80.-k,98.80.Es

I. INTRODUCTION

The past light cone, defined as the comoving distance r travelled by a light signal from time t to the present time $t_0\,(>t)$, in cosmology is given by [1]

$$r(t,t_0)=\frac{1}{H_0\sqrt{|\Omega_K|}}S_K\left(H_0\sqrt{|\Omega_K|}\int_t^{t_0}\frac{dt'}{a(t')}\right)\,, \qquad (1)$$

where $a(t)$ is the scale factor, H_0 is the Hubble parameter (defined as $H=d\ln a/dt$) computed at t_0 and $\Omega_K=-K/(H_0^2a_0^2)$ is the present curvature energy density ($K=1,0,-1$ for closed, flat, open spatial sections) with $a_0=a(t_0)$. Depending on $K=1\,,0\,,-1$ respectively, the function S_K:

$$S_k(x)=\begin{cases}\sin(x)\\ x\\ \sinh(x)\end{cases}$$

contains information on the geometry of space.

For flat spatial sections Eq. (1) simplifies to

$$r(t,t_0)=\int_t^{t_0}dt'/a(t')\,. \qquad (2)$$

Changing variable of integration from t to the redshift z (defined as $z=1/a-1$, with $a(t_0)=1$), one obtains

$$r(z)=\int_0^z dz'/H(z')\,. \qquad (3)$$

Since $H(z)$ is given by the Friedmann equation (see Sec.II), Eq. (3) allows the computation of r without solving for $a(t)$. Of course, once $r(z)$ is known, one still has to solve the Friedmann equation in order to obtain the dependence on time.

The integrations (2),(3) are the starting points for the computation of some distances of astrophysical interest as the Proper Distance (or Horizon), the Angular Diameter Distance and the Luminosity Distance [2]. The comoving distance (3) is also needed for the computation of the Acoustic Scale that is the characteristic angular scale of the peaks in the Power Spectrum of the Cosmic Microwave Background (henceforth CMB) anisotropies.

An analytic form for r is known for few simple cases [1] because is difficult to perform the integration either in form of Eq. (2) or Eq. (3). In the case of dust plus curvature, the time evolution for a and r are known [1].

Here we present the analytic solution of the scale factor *and* the exact computation of the past light cone in a flat Universe driven by Dust (whose energy density $\rho_C(t)\propto a^{-3}$) and Dark Energy ($\rho_X(t)\propto a^{\alpha}$) with a parameter of state (i.e. pressure over energy) constant in time. The important case of ΛCDM is obtained setting $\alpha=0$. We apply this formula to the following cases: first, we provide the exact expression of the Luminosity Distance (and we give an expansion of the past light cone at large z) and then we write down the analytic expression for the Acoustic Scale, relevant for the peaks analysis of the angular power spectrum of CMB anisotropies.

The paper is organized as follows: in section II we solve the Friedmann equation introducing a new time coordinate. In section III we find the analytic formula for the past light cone. We apply our results to the luminosity distance in section IV and to the acoustic scale in section V. We conclude in section VI.

II. SOLUTION OF THE FRIEDMANN EQUATION

In a flat universe, considering the FRW metric $ds^2=-dt^2+a^2(t)d\vec{x}^2$, filled with dust and another component with a constant parameter of state ($w_X=-1-\alpha/3$) we have

$$\rho_C(t)=\rho_C\,a^{-3}(t)\text{ and }\rho_X(t)=\rho_X\,a^{\alpha}(t)\,, \qquad (4)$$

where $\rho_X\,,\rho_C$ do not depend on time and are the energy density of the two components at present time. For $\alpha>-2$ the $X-$component drives the universe into asymptotic acceleration and for $\alpha>0$ such component

---

*Both the authors are also supported by INFN, Sezione di Bologna, via Irnerio, 46 – I-40126 Bologna – Italy

[†]Electronic address: gruppuso@iasfbo.inaf.it, finelli@iasfbo.inaf.it

violate the dominant energy condition ($w_X < -1$). The scale factor $a(t)$ satisfies the Friedmann equation

$$H^2 \equiv \left(\frac{da}{adt}\right)^2 = \frac{8\pi G}{3}\left[\rho_X a^\alpha + \frac{\rho_C}{a^3}\right] . \tag{5}$$

The total state parameter is

$$w_{\rm TOT}(a) = -\left(1+\frac{\alpha}{3}\right)\frac{\rho_X\, a^{\alpha+3}}{\rho_X\, a^{\alpha+3}+\rho_C} \tag{6}$$

and interpolates between 0 and w_X. It is not easy to find solutions for a two-component system (see however the radiation-matter universe and one of its generalizations [3]), but Eq. (5) can be integrated in the $\alpha = 0$ case (i.e. ΛCDM model) [4]. When $\alpha \neq 0$, it is no more possible to obtain analytically $a = a(t, \alpha)$ but only its inverse $t = t(a, \alpha)$. This problem can be solved defining a new time coordinate τ:

$$dt = d\tau\, a^{-\alpha/2} , \tag{7}$$

such that in this new coordinate, eq. (5) is given by

$$H_\tau^2 = \frac{8\pi G}{3}\left[\rho_X + \frac{\rho_C}{a^{3+\alpha}}\right] , \tag{8}$$

with $H_\tau = a'/a$, where the prime stands for derivative with respect to the time τ. Note that only for $\alpha = -2$, i.e. when the X-component behaves like a curvature term, τ becomes the conformal time η (defined as $d\eta = dt/a$). Eq. (8) can be integrated for $\alpha > -3$ with the result:

$$\begin{aligned} a(\tau) = &\left(\frac{\Omega_C}{1-\Omega_C}\right)^{1/(3+\alpha)} \times \\ &\times \left[\sinh\left(\frac{3+\alpha}{2}(1-\Omega_C)^{1/2}H_0\tau\right)\right]^{2/(3+\alpha)} , \end{aligned} \tag{9}$$

where $\Omega_C = 8\pi G\rho_C/3H_0^2 = 1-\Omega_X$ is the present dust density ratio [5]. For $\tau \to 0$, we find the standard dust behaviour $a \sim \tau^{2/(3+\alpha)} \sim t^{2/3}$. Notice that when $\alpha \to 0$, Eq. (9) recovers the known solution for ΛCDM model [4].

III. PAST LIGHT CONE

In order to perform the integration (2), we make use again of Eq. (7) so that is possible to rewrite the comoving distance as

$$r(a) = \int_a^1 \frac{da}{a^{1+\alpha/2}a'} . \tag{10}$$

By making explicit the form of H_τ in the integral, Eq. (10) reads

$$\begin{aligned} r(a) &= \left(\frac{1-\Omega_C}{\Omega_C}\right)^{\frac{(\alpha+2)}{2(\alpha+3)}} \frac{1}{H_0(1-\Omega_C)^{1/2}} \times \\ &\times \int_{a/A}^{1/A} \frac{dx}{x^{1/2}(1+x^{3+\alpha})^{1/2}} , \end{aligned} \tag{11}$$

where $1/A = (-1+\Omega_C^{-1})^{1/(3+\alpha)}$. The integration left in Eq. (11) can be performed giving (see 15.3.1 of [6])

$$\begin{aligned} r(z) &= \frac{1}{H_0}\frac{2}{\Omega_C^{1/2}}\left[f\left(\alpha, \frac{1}{\Omega_C}-1\right)\right. \\ &\quad \left. - \left(\frac{1}{1+z}\right)^{1/2} f\left(\alpha, \frac{1-\Omega_C}{\Omega_C(1+z)^{3+\alpha}}\right)\right] , \end{aligned} \tag{12}$$

where the definition of the redshift z has been used and where f is the following hypergeometric function [6]

$$\begin{aligned} f(\alpha, y) &= {}_2F_1\left[\frac{1}{2(\alpha+3)}, \frac{1}{2}, 1+\frac{1}{2(\alpha+3)}, -y\right] \\ &= \frac{(-y)^{\frac{-1}{(6+2\alpha)}}}{(6+2\alpha)} B_{-y}\left(\frac{1}{6+2\alpha}, \frac{1}{2}\right) , \end{aligned} \tag{13}$$

and in the last equality we have also rewritten the hypergeometric function in terms of incomplete Beta function $B_x(\gamma, \beta)$ [6].

It is interesting to expand the past light cone for large z, obtaining

$$\begin{aligned} r(z) = &\frac{2}{\Omega_C^{1/2}H_0}\left\{{}_2F_1\left[\frac{1}{2(\alpha+3)}, \frac{1}{2}, 1+\frac{1}{2(\alpha+3)},\right.\right. \\ &\left.\left. -\frac{1-\Omega_C}{\Omega_C}\right] - \frac{1}{(1+z)^{1/2}} + \mathcal{O}\left(\frac{1}{(1+z)^{\alpha+7/2}}\right)\right\} , \end{aligned} \tag{14}$$

where the leading term (that is constant with respect to the redshift z) is the conformal age of the universe already given in [7]. Note that the ΛCDM cosmic age in cosmic time is given in [1].

IV. LUMINOSITY DISTANCE

Eq. (12) is the starting point for the computation of distances of astrophysical interest such as the Luminosity and the Angular Distances. The Luminosity Distance, d_L, and the Angular Distance, d_A, in a flat universe are given by

$$d_L(z) = (1+z)\, r(z) , \tag{15}$$

$$d_A(z) = (1+z)^{-1}\, r(z) . \tag{16}$$

Let us focus on the Luminosity Distance. From Eq. (15), d_L can be expanded for small z by using its form in terms of incomplete Beta function

$$H_0 d_L(z) = z + d_2 z^2 + d_3 z^3 + d_4 z^4 + O(z^5) , \tag{17}$$

with

$$d_2 = \frac{(4+\alpha)-(3+\alpha)\Omega_C}{4} , \tag{18}$$



FIG. 1: Left Panel: Exact luminosity distance $H_0\,d_L$ (solid line) and approximated luminosity distance $H_0 d_{app}$ (up to third order, dotted line), both vs redshift z. Right Panel: p vs z where p is implicitly defined by $d_L = (1+p)d_{app}$ where d_L is the exact luminosity distance and d_{app} is the approximated (up to third order) luminosity distance. In both panels $\Omega_C = 0.3$ and $\alpha = 0$.

$$\begin{aligned} d_3 &= \frac{1}{24}\left[\alpha(4+\alpha) - 2(3+\alpha)(5+2\alpha)\Omega_C + \right. \\ &\quad \left. +3(3+\alpha)^2\Omega_C^2\right] \,, \end{aligned} \tag{19}$$

$$\begin{aligned} d_4 &= \frac{1}{192}\left[(-2+\alpha)\alpha(4+\alpha) \right. \\ &\quad -(3+\alpha)(40+\alpha(50+13\alpha))\Omega_C \\ &\quad \left. +3(3+\alpha)^2(20+9\alpha)\Omega_C^2 - 15(3+\alpha)^3\Omega_C^3\right] \end{aligned} \tag{20}$$

Once α is set to zero, d_2 and d_3 recover the known coefficients in flat ΛCDM model [8] with the deceleration parameter $q_0 = \Omega_C/2 - \Omega_X = 3\Omega_C/2 - 1$ and with the jerk parameter given by $j_0 = \Omega_C + \Omega_X = 1$.

In Fig. 1 we show the comparison between the exact (see Eq. (15)) and the approximated (up to the third order, see Eq. (17)) Luminosity Distance as a function of z for $\Omega_C = 0.3$ and $\alpha = 0$. For istance at $z = 1.5$, an error of 9.8% is made, if one considers the approximated instead of the exact luminosity distance.

In Fig. 2 we compare the exact luminosity distance with the first three approximated luminosity distances in a flat ΛCDM model. We note that the approximation of order z^2 always overestimate the exact luminosity distance while the approximations of order z^3 and z^4 always underestimate the exact luminosity distance. From this analysis it is clear that already for $z \sim 2$ the use of a third order expansion may flaw the comparison of observations with theoretical predictions.

In the left panel of Fig. 3 we set $\Omega_C = 0.3$ (and hence $\Omega_X = 0.7$) and plot Eq. (15) for different values of α. We notice that the dependence on α is quite weak. In the right panel of Fig. 3 we set $\alpha = 0$ (i.e. ΛCDM model) and plot Eq. (15) for different values of the ratio Ω_X/Ω_C. Of course larger is this ratio and larger is the luminosity distance for each fixed redshift. We notice that the dependence on this ratio is stronger than the α dependence.

V. ACOUSTIC SCALE

The characteristic angular scale of the peaks of the angular power spectrum in CMB anisotropies is defined [10] as

$$\theta_A = \frac{\pi}{\ell_A} = \frac{r_s(z_{dec})}{r(z_{dec})} \,, \tag{21}$$

where $r_s(z_{dec})$ is the comoving size of the sound horizon at decoupling, $r(z_{dec})$ is the comoving distance at decoupling given in Eq. (12) and ℓ_A is the multipole associated to the angular scale θ_A, also called acoustic scale. In fact, the expression (12) misses the contribution coming from the *radiation* (Ω_{rad}). A numerical study shows that we get an overestimate for ℓ_A that depends on α and in the worst case it is of the order of 0.5%. The comoving size of the sound horizon is given by

$$r_s(z_{dec}) = \int_0^{1/(1+z_{dec})} \frac{da}{a^2 H(a)}\, c_s(a) \,, \tag{22}$$

where c_s is the sound speed given by [11]

$$c_s(a) = 1\Big/\sqrt{3 + \frac{9}{4}\frac{\Omega_b}{\Omega_\gamma} a} \,, \tag{23}$$

where Ω_b and Ω_γ are respectively, the present density ratio for baryons and for photons. The integral (22) can



FIG. 2: Exact luminosity distance (solid line) and three approximated luminosity distance for $\Omega_C = 0.3$ and $\alpha = 0$ vs redshift. Long, short dashed and dotted line are second, third and fourth order in z expansion, respectively. Future cosmological interpretations of high-redshift data [9] need the accuracy of the exact solution.

be analytically performed once Ω_X, that enters in $H(a)$, is set to zero. The result is [11]

$$\begin{aligned} r_s(a_{dec}) &= \frac{4}{3H_0}\sqrt{\frac{\Omega_{rad}}{\Omega_C \Omega_b}} \times \\ &\times \ln\left[\frac{\sqrt{1+R_{dec}}+\sqrt{R_{dec}+R_{eq}}}{1+\sqrt{R_{eq}}}\right] \end{aligned}, \quad (24)$$

with $R(a) = 3(\Omega_b/(4\Omega_\gamma))a$ and where the label *dec* stands for "computed at decoupling" while *eq* stands for "computed at equivalence"(between radiation and matter). A numerical study shows that in the worst case this approximation (i.e. $\Omega_X = 0$ in r_s) gives an error of the order of $10^{-5}\,\%$, then completely negligible.

We are now ready to make a prediction for the acoustic scale ℓ_A defined in Eq. (21). We consider $\alpha = 0$ (i.e. ΛCDM) and set $a_{dec} = 1/1089$, $\Omega_C = 0.27$, $\Omega_b = 0.046$ and $\Omega_X = 0.73$. In order to compute z_{eq} we made use of the following relation [10]

$$1 + z_{eq} = \frac{5464\,(\omega_m/0.135)}{(T_{CMB}/2.725)^4(1+\rho_\nu/\rho_\gamma)}\,, \quad (25)$$

where $\omega_m = \omega_b + \omega_{cdm} = \Omega_C h^2$ (with $h = H_0/(100\,\mathrm{km}\,s^{-1}\,\mathrm{Mpc}^{-1})$, ρ_ν/ρ_γ is the ratio of neutrinos to photons densities and T_{CMB} is the CMB temperature in ${}^0K$. Setting $T_{CMB} = 2.725$, $h = 0.72$, $\rho_\nu/\rho_\gamma = 0.6851$ [10] we get $z_{eq} = 3362$. Once z_{eq} is known, it is possible to compute Ω_γ through

$$\Omega_\gamma = \frac{\Omega_C}{(1+\rho_\nu/\rho_\gamma)(z_{eq}+1)}\,, \quad (26)$$

obtaining $\Omega_\gamma = 4.76\;10^{-5}$. Thus we have all we need to compute the acoustic scale defined in Eq. (21). We obtain

$$\ell_A = 301.76\,, \quad (27)$$

which is in agreement with the Wilkinson Microwave Anisotropy Probe (WMAP) 1σ result [10]: $\ell_A^{WMAP} = 300 \pm 3$.

So our analytic calculation of the acoustic scale agrees with the value reported by the WMAP team. This is a further application of our results to a different set of observations, i.e. CMB anisotropies, which require a correct evaluation of the past light cone at high redshift, where a perturbative expansion does not work and only numerical methods were used so far.

Therefore, by using Eq. (21) and Eq. (14), the acoustic scale is given by:

$$\ell_A \simeq \frac{3\pi}{2}\sqrt{\frac{\Omega_b}{\Omega_\gamma}}\,\frac{{}_2F_1\left[-1/(6\omega_X), 1/2, 1-1/(6\omega_X), -(1-\Omega_C)/\Omega_C\right] - 1/(1+z_{dec})^{1/2}}{\ln\left[\sqrt{1+R_{dec}}+\sqrt{R_{dec}+R_{eq}}\right] - \ln\left[1+\sqrt{R_{eq}}\right]}\,, \quad (28)$$

where the dependence on the cosmological parameters and in particular on ω_X is explicit. Our Eq. (28) is completely analytic compared to $\Delta\ell$ given in [12], in which $r(z_{dec})$ contains a function F that is computed numerically [13] and $r_s(z_{dec})$ contains an averaged speed of sound that is fixed to a particular numerical value. In Ref. [12] the dark energy parameter of state is a function of time, but is taken as a time-average for the final expression of $\Delta\ell$ in the models considered in [12]. In Fig. 4 we show the dependence of the acoustic scale on α, keeping the other parameters fixed to the values reported above in the text.

VI. CONCLUSION

We have found the analytic solution for the scale factor *and* past light cone for a flat universe driven by dust and another X component with a parameter of state constant in time which dominates in the future. If $w_X < -1/3$, X is the simplest possible parametrization for Dark Energy. While Eq. (9) generalizes the previously known solution for ΛCDM [4] to $\alpha \neq 0$, the result (12) extends [7] for finite z, which is important for accurate quantitative predictions.



FIG. 3: Left Panel: Exact luminosity distance vs redshift for $\Omega_C = 0.3$ and $\alpha = -2$ (solid line), $\alpha = -1$ (dotted line), $\alpha = 0$ (short dashed), $\alpha = 1$ (dashed line) and $\alpha = 2$ (long dashed line). Right Panel: Exact luminosity distance vs redshift for $\alpha = 0$ and for $\Omega_X/\Omega_C = 9$ (solid line), $\Omega_X/\Omega_C = 7/3$ (dotted line), $\Omega_X/\Omega_C = 1$ (short dashed line), $\Omega_X/\Omega_C = 3/7$ (dashed line), $\Omega_X/\Omega_C = 1/9$ (long dashed line).



FIG. 4: Acoustic Scale vs α. Our analytic evaluation of the acoustic scale lies between the dashed lines. The higher dashed curve is computed with Eq. (21) while the lower is the 99.5% of the previous one. In this way we give an estimate of the order of magnitude of the error it is made, neglecting the contribution of the radiation in Eq. (12). Horizontal lines represent the 1σ contour to the value given by [10].

Eq. (12) has been applied to the following cases: the computation of the luminosity distance, relevant for supernovae, and the acoustic scale, which determines the position of the peaks in the pattern of CMB anisotropies. Both of these results are original. For the second case, our analytic prediction is in agreement with the value reported by the WMAP team. Note that this latter application requires a correct evaluation of the past light cone at high redshift, where a perturbative expansion does not work and only numerical methods were used so far.

A parameter of state constant in time cover a wide range of cosmological models which include dust. From a fundamental physics point of view, X corresponds to a cosmic string network or a curvature term when $\alpha = -2$ or to a domain walls background when $\alpha = -1$. A quintessence model with an exponential potential which does not have tracking behaviour leads to a component with $-2 < \alpha < 0$ (for $\alpha > 0$ one has to reverse sign of the kinetic term of a canonical scalar field, i.e. consider a phantom with an exponential potential).

[1] Kolb E. W., Turner M. S., 1989, *The Early Universe*, Addison-Wesley Publ. Co.

[2] Peebles P. J. E., 1971, *Physical Cosmology*, Princeton University Press

[3] Finelli F., 2003, JCAP 0310, 011

[4] Kofman L., Starobinsky, A. A., 1985, Sov. Astron. Lett. **11**, 271

---

[5] In analogy this trick can be used to find the solution for a bouncing universe by replacing ρ_X with $-\rho_X$, generalizing the matter plus positive curvature solution [1]. In terms of the new time τ the solution for the scale factor is:

$$a(\tau) = \left(\frac{\rho_C}{\rho_X}\right)^{1/(3+\alpha)} \left[\sin\left(\frac{3+\alpha}{2}\sqrt{\frac{8\pi G}{3}}\rho_X^{1/2}\tau + \varphi\right)\right]^{2/(3+\alpha)} ,$$

where φ is taken 0 for $\alpha > -3$ and $\pi/2$ for $\alpha < -3$. In this way the bounce is always at $\tau = 0$. Note that the allowed values for α are $-3 + 1/n$ with n being an non-vanishing integer.

[6] Abramowitz M. and Stegun I., *Handbook of Mathematical Functions*, 1985, (Dover Publ., New York)

[7] Doran M., Schwindt J. and Wetterich C., 2001, Phys.Rev.D **64**:123520

[8] V.Sahni, et al., 2003, JETP Lett. **77**, 201; Visser M., 2004, Class.Quant.Grav.**21**:2603-2616

[9] D. Hooper and S. Dodelson, arXiv:astro-ph/0512232.

[10] Page L., et al., 2003, ApJS **148**, 233

[11] Hu W. and Sugiyama N., 1995, Astrophys.J. **444**:489-506

[12] Doran M., Lilley M. J., Schwindt J. and Wetterich C., 2001, Astrophys. J. **559** 501

[13] The function F becomes the hypergeometric function given in eq. (14) when the contribution coming from radiation is neglected.